



**INTERNATIONAL JOURNAL OF  
PHARMACEUTICAL SCIENCES**  
[ISSN: 0975-4725; CODEN(USA): IJPS00]  
Journal Homepage: <https://www.ijpsjournal.com>



## Research Article

# Spectrophotometric Methods For Quantitative Estimation Of Tramadol Hydrochloride From Tablet Formulation

**Akash Ingale\***, Dr. Parag Patil, Vishwas More, Suraj Patil, Rahulsingh khairnar, Tushar Toke, Devendra Talele, Divyank Patil

*KYDSCTS College of Pharmacy Sakegaon Bhusawal.*

### ARTICLE INFO

Received: 11 July 2024

Accepted: 13 July 2024

Published: 19 July 2024

#### Keywords:

Tramadol Hydrochloride, Chloroform extract, Absorbance, Metanil yellow and Tropaeolin.

#### DOI:

10.5281/zenodo.12780200

### ABSTRACT

There are now two straightforward and highly accurate visible spectrophotometric techniques available for quantifying Tramadol hydrochloride from tablet form. The created techniques are predicated on the creation of a drug's colored, chloroform-extractable complex containing tropaeolin ooo and metanil yellow. Method I of the drug's chloroform extracted complex revealed an absorbance maxima at 410.0 nm and linearity in the concentration range of 324 µg/ml; method II of the drug's complex revealed an absorbance maxima at 485.0 nm and linearity in the concentration range of 324 µg/ml. The analysis results for the two created approaches were confirmed using statistical validation and recovery experiments.

### INTRODUCTION

Tramadol<sup>1</sup> Chemically it is [2-(dimethylaminomethyl)-1-(3-methoxyphenyl)cyclohexanol] hydrochloride. Tramadol hydrochloride is a centrally acting analgesic, used for treating moderate to severe pain. It possesses agonist actions at the µ-opioid receptor and effects reuptake at the noradrenergic and serotonergic systems. Tramadol is a compound with µ-agonist activity<sup>2</sup>. It is used to treat moderate to moderately severe pain and most types of neuralgia, including trigeminal neuralgia. Literature survey reveals that various methods available for estimation of

tramadol hydrochloride like UV spectrophotometric<sup>3-8</sup>, RP-HPLC<sup>9</sup>, LC-MS<sup>10</sup>, HPTLC<sup>11</sup>, etc. have been reported. However, there is no work in the literature reported about the visible Spectrophotometric estimation of Tramadol Hydrochloride in pharmaceutical dosage forms with Metanil yellow & Tropaeolin ooo reagents. An attempt has been made in the present study to develop two simple & rapid visible Spectrophotometric methods for analysis of Tramadol Hydrochloride in pharmaceutical dosage forms.

**\*Corresponding Author:** Akash Ingale

**Address:** KYDSCTS College of Pharmacy Sakegaon Bhusawal

**Email** ✉: [ingaleakash111@gmail.com](mailto:ingaleakash111@gmail.com)

**Relevant conflicts of interest/financial disclosures:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.



## Experimental

A Shimadzu UV/Visible spectrophotometer- 1700 with 1 cm matched quartz cells was used for spectral measurement. All the chemicals used were of analytical grade, metanil yellow solution (0.05% w/v) and tropaeolin ooo (0.03% w/v) was prepared in acid buffer<sup>9</sup> pH 1.6 and extracted several times with chloroform so as to remove chloroform soluble impurities. Buffer solutions were prepared in double distilled water. The tablet samples of Tramadol Hydrochloride were procured from the local market. For method I, in a series of 10 ml volumetric flasks aliquots of standard drug solution of Tramadol Hydrochloride (30  $\mu$ g/ml) in chloroform were transferred and diluted with the same so as to give several dilutions in the concentration range of 3-24  $\mu$ g/ml of drug. To 10 ml of each dilution taken in a separating funnel, 10 ml of metanil yellow solution was added, shaken and allowed to stand for 10 minutes for the formation of colored complex. The colored chloroform layer was separated out and absorbance was measured at 410.0 nm against a reagent blank. A calibration curve was prepared by plotting concentration versus absorbance. For method II, in a series of 10 ml volumetric flasks aliquots of standard drug solution of Tramadol Hydrochloride (30  $\mu$ g/ml) in chloroform were transferred and diluted with the same so as to give several dilutions in the concentration range of 3-24  $\mu$ g/ml of drug. To 10 ml of each dilution taken in a separating funnel, 10 ml of tropaeolin ooo solution was added, shaken and allowed to stand for 10 minutes for the formation of colored complex. The colored chloroform layer was separated out and absorbance was measured at 485.0 nm against a reagent blank. A calibration curve was prepared by plotting concentration versus absorbance.

For analysis of formulation, twenty tablets of Tramadol Hydrochloride were accurately weighed and average weight per tablet was determined. The

tablets were powdered and powder equivalent to 100 mg of Tramadol Hydrochloride was accurately weighed and extracted four times with 20 ml portions of chloroform, the combined extract was filtered through Whatmann filter paper No.41 into 100 ml volumetric flask. The residue was washed with chloroform and the washings were added to the filtrate, final volume of filtrate was made up to the mark with chloroform. From the above filtrate 3 ml was further diluted to 100 ml in a volumetric flask to get a tablet sample stock of 30  $\mu$ g/ml. For method I, 4 ml of above stock was further diluted to 10 ml with chloroform and for method II, 4 ml of above stock was further diluted to 10 ml with chloroform. This was treated as per the respective procedure for the calibration curve and absorbance was measured at 410.0 and 485.0 nm respectively and the amount of drug present in sample was computed from respective calibration curve.

## RESULTS AND DISCUSSION

Both the developed methods were repeated five times for two different strength of tablet formulation. Results of analysis are reported in Table 2. Recovery studies were carried out for both the developed methods by addition of known quantity of pure drug solution to pre analyzed tablet sample solution at three different concentration levels. The result of recovery studies is reported in Table 2. The proposed Spectrophotometric methods for determination of Tramadol Hydrochloride from tablet formulations are based on formation of chloroform extractable colored complex of drug with metanil yellow and tropaeolin ooo. The pH required for method I and II was optimized at pH 1.6 . The optical characteristics such as Beer's law limits, sandell's sensitivity, molar extinction coefficient and percent relative standard deviation, (calculated from the eight measurements with in the Beer's law limits) were calculated and the results are summarized in Table-1.



**Table-I Optical Characteristics And Precision Of Proposed Methods I And II**

Parameter	Method-I	Method-II
$\lambda_{max}$ (nm)	410.0	485.0
Beer's law limits ( $\mu\text{g/ml}$ )	3-24	3-24
Molar absorptivity ( $\text{Lmole}^{-1}\text{cm}^{-1}$ )	$0.1523 \times 10^6$	$0.1712 \times 10^6$
Sandell's sensitivity ( $\mu\text{g cm}^{-2}/0.001$ absorbance unit)	0.00172	0.00175
<b>Regression equation (<math>Y=a+b C</math>)</b>		
Slope (b)	0.0507	0.0574
Intercept (a)	-0.002	-0.0012
Correlation coefficient (r)	1.0000	1.0000
Relative standard deviation (%) *	0.3106	0.05964
<b>% Range of error (confidence limits) *</b>		
0.05 level	0.2597	0.0499
0.01 level	0.3842	0.0738
<b><math>Y = a + b C</math>, where C is concentration in <math>\mu\text{g/ml}</math> and Y is absorbance unit.</b>		
<b>* Eight replicate samples</b>		

**Table - II Assay And Recovery Of Tramadol Hydrochloride In Dosage Forms**

Sample	Labeled amount	Amount obtained (mg/tab) (mg/tab)		% Recovery**		
		Proposed method		Reference* method		
		I	II	I	II	
1	50	49.96	49.92	49.94	99.92	99.84
2	50	49.88	49.94	49.96	99.76	99.92
3	50	49.95	49.96	49.92	99.90	99.84
<b>* Reference U.V. Method developed in our lab</b>				<b>** Average of three determination</b>		

Regression characteristics like standard deviation of slope ( $S_b$ ), standard deviation of intercept ( $S_a$ ), standard error of estimation ( $S_c$ ), and percentage ranges of error (0.05 and 0.01 confidence limits) were calculated and are shown in Table-1.

Recovery studies were satisfactory which shows that there is no interference of excipients. The developed methods were found to be simple, rapid and accurate and can be used for routine analysis of drug from tablet formulations.

## REFERENCE

- Budvari S, Eds., In. 2001, The Merck Index, 13th Edn., Merck and Co., Inc., White house Station, NJ, 1706.
- Tripathi K.D. 2003, Essentials of Medical Pharmacology. New Delhi, Jaypee Publishers (P) Ltd, 5th Ed., 48-52.
- Garvendra S R, Pawan K B, Manish S, Roop N G. Estimation of Tramadol Hydrochloride from tablet formulation. Asian J. Chemistry, 2009; Vol. 21(8): 6111- 6115.
- Nagaraja KS, Ramachar T, Chakravarthy IE, Prabhavathi K. Simple Spectrophotometric estimation of Tramadol hydrochloride in pharmaceutical formulations. Chem. Sci. Trans., 2012; 1(2): 317-320.
- Lavanya G, Sujana K. Development and Validation of UV Spectroscopic method for determination of Tramadol hydrochloride in bulk & formulation. Int. J. Pharmacy & Ph.Sc., 2012; vol.4. suppl5:275-279.
- Sagarbechara, Padmavathi P. Prabhu, Subrahmanyam E.V.S. Development of new analytical method & its validation for the determination of tramadol hydrochloride in bulk & marketed formulations. Asian J. Biomedical & Ph. Sc, 2013; vol.3(23):53-56.



7. Trupti.P.Patel, Arun.M.Prajapati. Spectrophotometric estimation of Tramadol HCl & Aceclofenac in combined dosage form by second order derivative method. *Am. J. Pharm Tech Res.*, 2013;Vol.3(1): 999-1009.
8. Deepali G, Pandurang D. Simultaneous estimation of Tramadol hydrochloride and Paracetamol by UV Spectrophotometric method from tablet formulation. *Int.J.Pharm. Tech Research*,2010;vol.2(2): 1119-1123.
9. Sinha Manish, Verma Vivek. RP-HPLC method development & validation of Tramadol HCl in bulk form by Ion pair Liquid Chromatography. *J. Drug Delivery & Therapeutics*,2014;4(1):63-65.
10. Liu P, Liang, Wang BJ, Guo RC. Development & validation of a LC-MS method for the estimation of Tramadol in human plasma and urine. *Eur. J. Drug Metab. Pharmacokinetics*,2009;vol 34(3- 4):185-192.
11. P.Desai, A.Captain, S.Kamdar. Development & validation of HPTLC method for the estimation of Tramadol HCl in bulk & in capsule dosage form. *Int. J. Pharm. Tech. Research*, 2012; vol.4(3):1261-1265.
12. 1996, Indian pharmacopoeia,4th Edn., Vol. II, A-144.

**HOW TO CITE:** Akash Ingale\*, Dr. Parag Patil, Vishwas More, Suraj Patil, Rahulsingh khairnar, Tushar Toke, Devendra Talele, Divyank Patil, Spectrophotometric Methods For Quantitative Estimation Of Tramadol Hydrochloride From Tablet Formulation, *Int. J. of Pharm. Sci.*, 2024, Vol 2, Issue 7, 1430-1433. <https://doi.org/10.5281/zenodo.12780200>